JEE Main April 2025

Question Paper With Text Solution

04 April | Shift-2

CHEMISTRY

MATRIX

JEE Main & Advanced | XI-XII Foundation| VI-X Pre-Foundation

Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911

Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in

JEE MAIN APRIL 2025 | 04TH APRIL SHIFT-2

SECTION – A

51. A dipeptide, "x" on complete hydrolysis gives "y" and "z". "y" on treatment with aq. HNO_2 produces lactic acid. On the other hand "z" on heating gives the following cyclic molecule.



Based on the information given, the dipeptide X is :

(1) alanine-alanine

(2) valine-glycine

(3) valine-leucine

(4) alanine-glycine

603421820

Ans. Official answer NTA (4)

Sol.

52. Which among the following compounds give yellow solid when reacted with $NaOI / NaOH$

(A) 

(B) $CH_3 - CH_2 - CH_2 - OH$

(C) 

(D) 

(E) 

Choose the correct answer from the options given below :

(1) (B), (C) and (E) Only

(2) (A), (C) and (D) Only

(3) (A) and (C) Only

(4) (C) and (D) Only

603421817

Ans. Official answer NTA (3)

Sol.

53. Consider the ground state of chromium atom (Z=24). How many electrons are with

Azimuthal quantum number l=1 and l=2 respectively ?

(1) 16 and 4

(2) 12 and 4

(3) 16 and 5

(4) 12 and 5

603421801

Ans. Official answer NTA (4)

Sol.

54. Given below are two statements :

Statement (I) : Alcohols are formed when alkyl chlorides are treated with aqueous potassium hydroxide by elimination reaction.

Statement (II) : In alcoholic potassium hydroxide, alkyl chlorides form alkenes by abstracting the hydrogen from the β-carbon.

In the light of the above statements, choose the most appropriate answer from the options given below :

(1) Statement I is incorrect but Statement II is correct

(2) Statement I is correct but Statement II is incorrect

(3) Both Statement I and Statement II are incorrect

(4) Both Statement I and Statement II are correct

603421816

Ans. Official answer NTA (1)

Sol.

55. Consider the following molecule (X).

The structure of X is



The major product formed when the given molecule (X) is treated with HBr(1eq) is :

(1) 

(2) 

(3) 

(4) 

603421815

Ans. Official answer NTA (3)

Sol.

56. Given below are two statements :

Statement (I) : For $\overset{\cdot\cdot}{\underset{\cdot\cdot}{C}}IF_3$, all three possible structures may be drawn as follows.



I



II



III

Statement (II) : Structure III is most stable, as the orbitals having the lone pairs are axial, where the lp – bp repulsion is minimum.

In the light of the above statements, choose the most appropriate answer from the options given below :

(1) Statement I is incorrect but Statement II is correct

(2) Both Statement I and Statement II are incorrect

(3) Both Statement I and Statement II are correct

(4) Statement I is correct but Statement II is incorrect

603421806

Ans. Official answer NTA (4)

Sol.

57. The elements of Group 13 with highest and lowest first ionisation enthalpies are respectively:

(1) B & Ga

(2) Tl & B

(3) B & Tl

(4) B & In

603421807

Ans. Official answer NTA (4)

Sol.

58. The correct order of $[FeF_6]^{3-}$, $[CoF_6]^{3-}$, $[Ni(CO)_4]$ and $[Ni(CN)_4]^{2-}$ complex species based on the number of unpaired electrons present is :

(1) $[CoF_6]^{3-} > [FeF_6]^{3-} > [Ni(CO)_4] > [Ni(CN)_4]^{2-}$

(2) $[FeF_6]^{3-} > [CoF_6]^{3-} > [Ni(CN)_4]^{2-} > [Ni(CO)_4]$

(3) $[Ni(CN)_4]^{2-} > [FeF_6]^{3-} > [CoF_6]^{3-} > [Ni(CO)_4]$

(4) $[FeF_6]^{3-} > [CoF_6]^{3-} > [Ni(CN)_4]^{2-} = [Ni(CO)_4]$

603421810

Ans. Official answer NTA (4)

Sol.

59. The incorrect relationship in the following pairs in relation to ionisation enthalpies is :

(1) $Fe^{2+} < Fe^{3+}$

(2) $Mn^{2+} < Fe^{2+}$

(3) $Mn^{+} < Cr^{+}$

(4) $Mn^{+} < Mn^{2+}$

603421811

Ans. Official answer NTA (2)

Sol.

60. Given below are two statements :

Statement (I) : Molal depression constant K_f is given by $\frac{M_1RT_f}{\Delta S_{fus}}$, where symbols have their usual meaning.

Statement (II) : K_f for benzene is less than the K_f for water.

In the light of the above statements, choose the most appropriate answer from the options given below :

(1) Statement I is incorrect but Statement II is correct

(2) Statement I is correct but Statement II is incorrect

(3) Both Statement I and Statement II are incorrect

(4) Both Statement I and Statement II are correct

603421803

Ans. Official answer NTA (2)

61. 'X' is the number of electrons in t_{2g} orbitals of the most stable complex ion among $\left[Fe(NH_3)_6\right]^{3+}$, $[FeCl_6]^{3-}$, $\left[Fe(C_2O_4)_3\right]^{3-}$ and $\left[Fe(H_2O)_6\right]^{3+}$. The nature of oxide of vanadium of the type V_2O_X is :

(1) Amphoteric

(2) Basic

(3) Neutral

(4) Acidic

603421809

Ans. Official answer NTA (1)

Sol.

62. Match List - I with List - II.

List - I (Separation of)	List - II (Separation Technique)
(A) Aniline from aniline-water mixture	(I) Simple distillation
(B) Glycerol from spent-lye in soap industry	(II) Fractional distillation
(C) Different fractions of crude oil in petroleum industry	(III) Distillation at reduced pressure
(D) Chloroform-Aniline mixture	(IV) Steam distillation

Choose the correct answer from the options given below :

(1) (A)-(IV), (B)-(III), (C)-(II), (D)-(I)

(2) (A)-(II), (B)-(I), (C)-(IV), (D)-(III)

(3) (A) – (III), (B) – (IV), (C) – (I), (D) – (II)

(4) (A) – (I), (B) – (II), (C) – (III), (D) – (IV)

603421812

Ans. Official answer NTA (1)

Sol.

63. In which pairs, the first ion is more stable than the second?

(A) 

(B) 

(C) 

(D) 

(1) A and B only

(2) B and C only

(3) B and D only

(4) A and C only

603421813

Ans. Official answer NTA (1)

Sol.

64. Consider the given data :

(a) $HCl(g) + 10H_2O(l) \rightarrow HCl.10H_2O \Delta H = -69.01 kJ\,mol^{-1}$

(b) $HCl(g) + 40H_2O(l) \rightarrow HCl.40H_2O \Delta H = -72.79 kJ\,mol^{-1}$

Choose the correct statement :

(1) The heat of solution depends on the amount of solvent.

(2) Dissolution of gas in water is an endothermic process.

(3) The heat of formation of HCl solution is represented by both (a) and (b).

(4) The heat of dilution for the $HCl\left(HCl.10H_2O \text{ to } HCl.40H_2O\right)$ is $3.78\ kJ\,mol^{-1}$.

603421802

Ans. Official answer NTA (1)

Sol.

65. The IUPAC name of the following compound is :



(1) Hept-6-en-1-yn-4-ol

(2) 4-Hydroxyhept-1-en-6-yne

(3) 4-Hydroxyhept-6-en-1-yne

(4) Hept-1-en-6-yn-4-ol

603421814

Ans. Official answer NTA (4)

Sol.

66. Given below are two statements :

Statement (I): The first ionisation enthalpy of group 14 elements is higher than the corresponding elements of group 13.

Statement (II) : Melting points and boiling points of group 13 elements are in general much higher than those of corresponding elements of group 14.

In the light of the above statements, choose the most appropriate answer from the options given below :

(1) Both Statement I and Statement II are correct

(2) Both Statement I and Statement II are incorrect

(3) Statement I is incorrect but Statement II is correct

(4) Statement I is correct but Statement II is incorrect

603421808

Ans. Official answer NTA (4)

Sol.

67. The correct order of basicity for the following molecules is :


(P)


(Q)


(R)

(1) Q > P > R

(2) P > Q > R

(3) R > P > Q

(4) R > Q > P

603421819

Ans. Official answer NTA (4)

Sol.

68. A toxic compound "A" when reacted with NaCN in aqueous acidic medium yields an edible cooking component and food preservative "B". "B" is converted to "C" by diborane and can be used as an additive to petrol to reduce emission. "C" upon reaction with oleum at $140^{\circ}C$ yields an inhalable anesthetic "D". Identify "A", "B", "C" & "D", respectively :

(1) Ethanol; acetonitrile; ethylamine; ethylene

(2) Methanol; formaldehyde; methyl chloride; chloroform

(3) Methanol; acetic acid; ethanol; diethyl ether

(4) Acetaldehyde; 2-hydroxypropanoic acid; propanoic acid; dipropyl ether

603421818

Ans. Official answer NTA (3)

Sol.

69. Consider the following plots of log of rate constant $k(\log k)$ vs $\frac{1}{T}$ for three different reactions. The correct order of activation energies of these reactions is :



(1) $Ea_1 > Ea_2 > Ea_3$

(2) $Ea_3 > Ea_2 > Ea_1$

(3) $Ea_1 > Ea_3 > Ea_2$

(4) $Ea_2 > Ea_1 > Ea_3$

603421805

Ans. Official answer NTA (4)

Sol.

70. Half life of zero order reaction $A \rightarrow$ product is 1 hour, when initial concentration of reactant is $2.0\,mol\,L^{-1}$. The time required to decrease concentration of A from 0.50 to $0.25\,mol\,L^{-1}$ is :

(1) 4 hour

(2) 15 min

(3) 0.5 hour

(4) 60 min

603421804

Ans. Official answer NTA (2)

Sol.

SECTION – B

71. The molar conductance of an infinitely dilute solution of ammonium chloride was found to be $185\,S\,cm^2\,mol^{-1}$ and the ionic conductance of hydroxyl and chloride ions are 170 and $70\,S\,cm^2\,mol^{-1}$, respectively. If molar conductance of 0.02 M solution of ammonium hydroxide is $85.5\,S\,cm^2\,mol^{-1}$, its degree of dissociation is given by $x\times10^{-1}$.

The value of x is ____. (Nearest integer)

603421822

Ans. Official answer NTA (3)

Sol.

72. Sea water, which can be considered as a 6 molar (6M) solution of NaCl, has a density of $2\,g\,mL^{-1}$. The concentration of dissolved oxygen (O_2) in sea water is 5.8 ppm. Then the concentration of dissolved oxygen (O_2) in sea water, is $x\times10^{-4}\,m$.

x= _____ (Nearest integer)

Given: Molar mass of NaCl is $58.5\,g\,mol^{-1}$

Molar mass of O_2 is $32\,g\,mol^{-1}$

603421823

Ans. Official answer NTA (2)

Sol.

73. A metal complex with a formula $MCl_4\cdot3NH_3$ is involved in sp^3d^2 hybridisation. It upon reaction with excess of $AgNO_3$ solution gives 'x' moles of AgCl. Consider 'x' is equal to the number of lone pairs of electron present in central atom of BrF_5. Then the number of geometrical isomers exhibited by the complex is _____.

603421825

Ans. Official answer NTA (2)

Sol.

74. The amount of calcium oxide produced on heating 150 kg limestone (75 % pure) is ___ kg. (Nearest integer)

Given: Molar mass (in g mol^{-1}) of Ca-40, O-16, C-12

603421824

Ans. Official answer NTA (63)

Sol.

75. 75xmg of $Mg(OH)_2$ (molar mass =58) is required to be dissolved in 1.0 L of water to produce a pH of 10.0 at 298 K . The value of x is _____ mg. (Nearest integer)

(Given : $Mg(OH)_2$ is assumed to dissociate completely in H_2O]

603421821

Ans. Official answer NTA (3)

Sol.